125<sub>T</sub>

Caution: For Laboratory Use. A product for research purposes only

## [<sup>125</sup>I]Tyr<sup>4</sup>-Angiotensin II (human) Asp-Arg-Val-[<sup>125</sup>I]Tyr-Ile-His-Pro-Phe

**Product Number: NEX105** 

## **LOT SPECIFIC INFORMATION:**

CALCULATED AS OF: 10-Feb-2014

LOT NUMBER: AL31440

SPECIFIC ACTIVITY: 81.4 TBq/mmol

2200 Ci/mmol 70 MBq/μg 1879 μCi/μg **Package Size Information** 

r donago oizo imormation
Package Size
as of
14-Mar-2014
370 kBq
10 μCi
1.85 MBq
50 μCi

**RADIOCHEMICAL PURITY:** ≥ 95%

MOLECULAR WEIGHT: 1,171

**PACKAGING**: [125]Tyr<sup>4</sup>-Angiotensin II is lyophilized from a solution containing 0.04M sodium phosphate, 1M glycine, 0.2M NaCl, 0.25% BSA, 500 KIU/ml Trasylol<sup>®</sup> at pH 7.2. It is shipped ambient.

**STABILITY AND STORAGE**: The lyophilized [ $^{125}$ I]Tyr $^4$ -Angiotensin II should be stored at  $4\,^{\circ}$ C or lower. Following reconstitution with distilled water to a concentration of approximately 50  $\mu$ Ci/ml on calibration date, aliquot and store at -20 $\,^{\circ}$ C or lower. Under these conditions the product is stable and usable in radioimmunoassays for at least eight weeks after fresh lot date.

**SPECIFIC ACTIVITY**: The initial specific activity of  $[^{125}I]Tyr^4$ -Angiotensin II is 2200 Ci/mmol (81 TBq/mmol), 1879  $\mu$ Ci/ $\mu$ g (70 MBq/ $\mu$ g). Preparative HPLC is used to separate unlabeled angiotensin II from  $[^{125}I]Tyr^4$ -Angiotensin II. Upon decay,  $[^{125}I]Tyr^4$ -Angiotensin II undergoes decay catastrophe and the specific activity remains constant with time. However, it is not known what molecular or peptide fragments are generated from the decay event or what functional activity these fragments may have in different assays. References on  $^{125}I$  decay and decay catastrophe of  $^{125}I$  labeled compounds are available.  $^{1-5}I$ 

RADIOCHEMICAL PURITY: Initially greater than 95% radiochemically pure as determined by HPLC.

**PREPARATIVE PROCEDURE**: Angiotensin II is radioiodinated with no carrier added <sup>125</sup>I using a modification of the Hunter and Greenwood method<sup>6</sup> and purified by reversed phase HPLC.

**AVAILABILITY:** [125] Tyr4-Angiotensin II is routinely available from stock and is prepared fresh and packaged for shipment on the second Monday of each month. Please inquire for larger package sizes.

**HAZARD WARNING**: This product contains a chemical (s) known to the state of California to cause cancer. This product also contains a component which is harmful by contact, ingestion and inhalation. It is irritating to the eyes, skin and respiratory tract and is toxic.

**RADIATION UNSHIELDED:** 280mR/hr/mCi at vial surface.

## **REFERENCES:**

- 1. Doyle, V.M., Buhler, F.R., Burgisser, E., Eur. J. Pharm. <u>99</u> 353 (1984).
- 2. Schmidt, J., J. Biol. Chem. 259 1660 (1984).
- 3. Loring, R.H., Jones, S.W., Matthews-Bellinger, J., Salpeter, M.M., J. Biol. Chem. 257 1418 (1982).
- 4. Berridge, M.S., Jiang, V.W., Welch, M.J., *Rad. Res.* <u>82</u> 467 (1980).
- 5. Charlton, D.E., Rad. Res. <u>107</u> 163 (1986).
- 6. Hunter, W.M. and Greenwood, F.C., *Nature*, <u>194</u>, 495 (1962).

## **IODINE-125 DECAY CHART HALF LIFE=60 days**

Radiations: Gamma 35.5 keV (7%), X-ray K alpha 27 KeV (112%), K beta 31 keV (24%)

DAYS	0	2	4	6	8	10	12	14	16	18
0	1.000	.977	.955	.933	.912	.891	.871	.851	.831	.812
20	.794	.776	.758	.741	.724	.707	.691	.675	.660	.645
40	.630	.616	.602	.588	.574	.561	.548	.536	.524	.512
60	.500	.489	.477	.467	.456	.445	.435	.425	.416	.406
80	.397	.388	.379	.370	.362	.354	.345	.338	.330	.322
100	.315	.308	.301	.294	.287	.281	.274	.268	.262	.256
120	.250	.244	.239	.233	.228	.223	.218	.213	.208	.203

To obtain the correct radioactive concentration or amount for a date before the calibration date: divide by the decay factor corresponding to the number of days before the calibration date. To obtain the correct radioactive concentration or amount for a date after the calibration date: multiply by the decay factor corresponding to the number of days after the calibration date.

PerkinElmer, Inc.

549 Albany Street

Boston, MA 02118 USA

P: (800) 762-4000 or (+1) 203-925-4602

www.perkinelmer.com/nenradiochemicals

For a complete listing of our global offices, visit

www.perkinelmer.com/ContactUs

Copyright ©2010, PerkinElmer, Inc. All rights reserved. PerkinElmer® is a registered trademark of PerkinElmer, Inc.

All other trademarks are the property of their respective owners.

